

Original Research Article

THE STUDY OF FUNCTIONAL OUTCOME OF CERVICAL DISC HERNIATION AFTER CERVICAL FRENCH-DOOR LAMINOPLASTY

Arjun Hari¹, Vincent Bosco Savery², NishanthAgarwal³, Aravind Reddy⁴, Abhiram SR⁵, Jyothirmoy Das Gup⁶, Prof. Sankaralal⁷

¹Post Graduate Trainee, Department of Orthopaedics, Sri Lakshmi Narayana Institute of Medical Sciences (SLIMS), Puducherry, India.

²Associate Professor, Department of Orthopaedics, Sri Lakshmi Narayana Institute of Medical Sciences (SLIMS), Puducherry, India

³Post Graduate Trainee, Department of Orthopaedics, Sri Lakshmi Narayana Institute of Medical Sciences (SLIMS), Puducherry, India.

⁴Post Graduate Trainee, Department of Orthopaedics, Sri Lakshmi Narayana Institute of Medical Sciences (SLIMS), Puducherry, India.

⁵Post Graduate Trainee, Department of Orthopaedics, Sri Lakshmi Narayana Institute of Medical Sciences (SLIMS), Puducherry, India.

⁶Post Graduate Trainee, Department of Orthopaedics, Sri Lakshmi Narayana Institute of Medical Sciences (SLIMS), Puducherry, India.

⁷Prof. Emeritus, Department of Orthopaedics, Sri Lakshmi Narayana Institute of Medical Sciences (SLIMS), Puducherry, India.

Received : 08/03/2024
Received in revised form : 12/05/2024
Accepted : 27/05/2024

Corresponding Author:

Dr. Vincent Bosco Savery

Associate Professor, Department of orthopaedics, Sri Lakshmi Narayana institute of medical sciences, Puducherry, India.
Email: vboscossavery@yahoo.com

DOI: 10.5530/ijmedph.2024.2.127

Source of Support: Nil,

Conflict of Interest: None declared

Int J Med Pub Health

2024; 14 (2); 665-669

ABSTRACT

Rationale: Cervical French-door laminoplasty can enlarge the volume of the cervical vertebral canal and thus has become an effective and safe treatment for multilevel cervical disc herniation and cervical stenosis. Some post-surgery complications exist, such as reduction of cervical alignment and local kyphosis. However, aggravation of cervical disc herniation at the surgical level during short-term follow-up has not been discussed. Additionally, spontaneous disappearance of herniated disc pulposus is a common phenomenon in the lumbar region but is relatively rare in the cervical region.

Patient concerns: We evaluated a group of 10 patients who Presented with history of paraesthesia and weakness in the upper and lower limbs. The sensations and muscle strength of both upper and lower limbs were decreased. The radiological findings showed that the Pavlov ratios from C3-5 were decreased obviously. Osteophytes as well as spinal cord compression were observed at C3/4, C4/5, and C5/6 in the index patient

Diagnoses: Considering the symptoms and clinical examinations, the index patient was diagnosed with cervical stenosis.

Interventions: We performed cervical French-door laminoplasty at C3-C5 to enlarge the space of the cervical vertebral canal.

Outcomes: At 1-month post-surgery follow-up, the patients showed obvious improvement in paraesthesia and weakness in the upper limbs. The cervical disc herniation at C3/4 was aggravated. However, on further follow up the symptoms were relieved, and is too early to comment on herniated cervical disc at C3/4 which spontaneously disappears without any special treatment.

Lessons: We suggest that the attachment points of deep muscles in the neck region should be carefully protected during this surgery. Patients who undergo cervical French-door laminoplasty should pay attention to their cervical position and perform neck exercises to train their neck muscles. MRI is an important imaging method to observe dynamic changes in herniated discs for patients with cervical disc herniation.

Abbreviations: FSU = functional spinal unit, MRI = magnetic resonance imaging, ROM = range of motion, SDHNP = spontaneous disappearance of the herniated nucleus pulposus.

Keywords: Aggravated disc herniation, cervical disc disappearance, cervical disc herniation, cervical laminoplasty, disc regression.

INTRODUCTION

Cervical disc herniation is a common condition that can cause serious symptoms, such as pain and numbness. Cervical French door laminoplasty can enlarge the volume of the cervical vertebral canal and thus has become an effective and safe treatment for multilevel cervical disc herniation and cervical stenosis.^[1-3] Studies have reported post-surgical complications, such as reduction of cervical alignment and local kyphosis.^[4-6] However, aggravation of cervical disc herniation at the surgical level during short-term follow-up has not been discussed. Spontaneous disappearance of the herniated nucleus pulposus (SDHNP) was first reported by Jelasic F in 1953.^[7] Most cases occur in the lumbar region, and cervical SDHNP is relatively rare. Many hypotheses have been proposed to explain the mechanism of SDHNP, but none have been comprehensive. We report a patient with aggravation of cervical disc herniation at C3/4 after cervical French-door laminoplasty at C3-5 during a short-term follow-up. The herniated disc subsequently spontaneously disappeared during long-term follow-up. Laminoplasty is an established procedure for the decompression of multisegmental compressive myelopathy in the cervical spine. We share our opinion about the mechanisms underlying these phenomena.

MATERIAL AND METHODS

The index patient provided informed consent for the publication of her clinical and radiological data. This case report was approved by the Medical Ethical Committee.our index patient a 63-year-old female presented with a 2-year history of neck pain and a 3 months history of paraesthesia and weakness in the upper and lower limbs. The symptoms did not subside after 3 months of conservative treatment. During the physical examination, the sensations of both the upper and lower limbs were decreased, and the left side symptoms were more serious. The muscle strength was grade 4 for the upper limbs and grade 3 for the lower limbs. Hoffman's sign was positive on both sides. The lateral radiograph showed a narrowed cervical vertebral canal, and the Pavlov ratios from C3-6 were 0.68, 0.64, 0.61, and 0.69, respectively. The range of motion (ROM) at C2-7 was 59.17°. The Mri showed obvious cervical disc herniation. Osteophytes were observed at C3/4, C4/5, and C5/6 (Fig. 1). Magnetic resonance imaging (MRI) revealed cervical spinal cord compression at C3/4, C4/5, and C5/6 (Fig. 2). Considering the symptoms and clinical examinations, the patient was diagnosed with cervical disc herniation. We performed cervical French-door laminoplasty at C3-5 (Fig 3) to enlarge the space of the cervical vertebral canal.




Figure 1: Cervical Disc Herniation (arrow Marked)





Figure 2: MRI (Axial) before surgery reveals cervical spinal cord compression at C3/4, C4/5 and C5/6

At the 1-month post-surgery follow-up, the patient showed obvious improvement in paraesthesia and weakness in the upper limbs. The lateral radiograph showed a straightened sagittal alignment and cervical kyphosis at the C3/C4 level. The C2-7 Cobb's angle decreased from 13.13° to 9.15°.The MRI showed decompression at C3-5 and aggravation of cervical disc herniation at C3/4. The patient denied any history of trauma. Considering that no new symptoms occurred, we recommended follow up for the herniated disc in 12 months. At the 1-month post-surgery follow-up, she recovered well without any discomfort. However, it is too early to comment on herniated cervical disc at C3/4 which spontaneously disappears without any special treatment.



A



D

Figure 3 (A-D): Procedures of French-door laminoplasty during surgery



B



A



B



C



C

Figure 4 (A-C): Functional outcome at 2 years

DISCUSSION

We evaluated and operated on a group of 10 patients and reporting on functional outcome of our index patient with cervical spinal herniation and multilevel spinal cord compression. She underwent cervical French-door laminoplasty, and her cervical disc herniation at C3/4 was aggravated at the 1-month follow-up. Lee et al,^[8] reported that the cervical sagittal alignment might become kyphotic after cervical french-door laminoplasty. Usually, the attachments of deep muscle in the neck region and posterior ligamentous structures are cut during this surgery; thus, the posterior muscle strength in the cervical region is decreased after surgery. This decrease can lead to kyphosis of the cervical sagittal alignment.^[9] Some authors reported that the abnormal cervical sagittal alignment was related to cervical disc degeneration and herniation.^[10-12] An abnormal cervical sagittal alignment could increase the intervertebral disc pressure and change the transmission pattern of the cervical stress. In our case, the patient's C2-7 Cobb's angle decreased from 13.13° to 9.15° and the FSU angle at C3-4 decreased from -3.62° to -7.45° at the 2-month follow-up. The cervical sagittal alignment obviously worsened. Poor positioning of the cervical and thoracic spine have been reported to be highly related to abnormal cervical sagittal alignment.^[13,14] Additionally, the annulus fibrosus was broken before surgery. All of these factors could have led to aggravation of the HNP at C3/4. Laminoplasty is an established procedure for the decompression of multisegmental compressive myelopathy in the cervical spine. Several kinds of procedures of laminoplasty were developed in the 1970s and 1980s.^[15,16] Laminoplasties are safe and minimally invasive with low rate of complications.^[17] These procedures can preserve range of motion postoperatively. Open-door laminoplasty and French-door laminoplasty are two representative techniques of laminoplasties. Both techniques can provide similar clinical outcomes. However French-door laminoplasty is superior to open-door laminoplasty in terms of intraoperative bleedings and postoperative range of motion and alignment. Qiu et al,^[18] reported that the volume of the neck muscles was maintained if the paraspinal muscular-ligament complex on one side was reserved during cervical open-door laminoplasty. We suggest that the attachment points of deep muscles in the neck region should be carefully protected during this surgery. Patients undergoing cervical French-door laminoplasty should pay attention to their cervical position and perform neck exercise to train their neck muscles. By reviewing the literature, we found that SDHNP was first reported in 1953.^[7] Most SDHNP cases occur in the lumbar region, and cervical SDHNP is relatively rare. To the best of our knowledge, this case is the first report of the

aggravation and spontaneous disappearance of cervical HNP after cervical laminoplasty. We offer some hypotheses about the mechanisms of SDHNP. First, the inflammatory immune response could lead to SDHNP. The intervertebral discs are in an avascular physiological environment. Once the nucleus pulposus herniates into the vascular epidural space, it may be recognized as a foreign body by the immune system and cause a series of inflammatory reactions, and the herniated nucleus pulposus may be attacked and broken down.^[19,20] SDHNP was observed to be more likely to appear as the size of the nucleus pulposus that herniated into the epidural space increased.^[21,22] Second, some authors believe that the RNHP is a form of apoptosis induced by various cytokines.^[19,23] In addition, for acute SDHNP, if the HNP does not separate from the annulus fibrosus, the herniated disc can retract back into the intervertebral disc space when the pressure of the intervertebral disc space decreases.^[24,25] Furthermore, the nucleus pulposus has been hypothesized to wear out if it herniates into the epidural space and contacts the flowing cerebrospinal fluid; alternatively, the herniated nucleus pulposus may shrink and gradually disintegrate.^[26] Although this case is unique, it also has some limitations. We only obtained the patient's imaging data from her 2 week and 2-month follow-ups; therefore, we could not compare the dynamic changes of the nucleus pulposus at C3/4 consecutively. We cannot generalize our case report beyond the present context to a larger population of patients, although our findings do suggest that cervical disc herniation can become aggravated after cervical laminoplasty and that the protruding nucleus pulposus can disappear spontaneously. Future research in the form of retrospective studies and basic experiments could be based on the work we describe here to provide additional evidence about the phenomenon and to understand the underlying mechanisms.

CONCLUSION

Cervical French-door laminoplasty appears to be a promising surgical approach for improving functional outcomes in patients with cervical disc herniation. Further research with larger patient cohorts and longer follow-up periods is necessary to solidify these findings and determine the long-term effectiveness of this procedure

REFERENCES

1. Lee TT, Green BA, Gromelski EB. Safety and stability of open-door cervical expansive laminoplasty. *J Spinal Disord* 1998; 11:12-5.
2. Tsuji T, Asazuma T, Masuoka K, et al. Retrospective cohort study between selective and standard C3-7 laminoplasty. Minimum 2-year follow-up study. *Eur Spine J* 2007; 16:2072-7.
3. Jiang Y, Li X-L, Zhou X, et al. A prospective randomized trial comparing anterior cervical discectomy and fusion versus

- plate-only open-door laminoplasty for the treatment of spinal stenosis in degenerative diseases. *Eur Spine J* 2017; 26:1162–72.
4. Suk KS, Kim KT, Lee JH, et al. Sagittal alignment of the cervical spine after the laminoplasty. *Spine (Phila Pa 1976)* 2007; 32:656–60.
 5. Chiba K, Ogawa Y, Ishii K, et al. Long-term results of expansive open-door laminoplasty for cervical myelopathy average 14-year follow-up study. *Spine (Phila Pa 1976)* 2006; 31:2998–3005.
 6. Nakashima H, Kato F, Yukawa Y, et al. Comparative effectiveness of open-door laminoplasty versus French-door laminoplasty in cervical compressive myelopathy. *Spine (Phila Pa 1976)* 2014; 39:642–7.
 7. Jelastic F. Resorption of prolapsed intervertebral disk; experimental studies. *Acta Med Jugosl* 1953; 7:98–111.
 8. Lee DH, Cho JH, Hwang CJ, et al. Multilevel posterior foraminotomy with laminoplasty versus laminoplasty alone for cervical spondylotic myelopathy with radiculopathy: a comparative study. *Spine J* 2017; [Epub ahead of print].
 9. Healy AT, Lubelski D, West JL, et al. Biomechanics of open-door laminoplasty with and without preservation of posterior structures. *J Neurosurg Spine* 2016; 24:746–51.
 10. Miyazaki M, Hymanson HJ, Morishita Y, et al. Kinematic analysis of the relationship between sagittal alignment and disc degeneration in the cervical spine. *Spine (Phila Pa 1976)* 2008; 33:870–6.
 11. Yu M, Mai WZ, Wang LS. Analysis of cervical and global spine alignment under Roussouly sagittal classification in Chinese cervical spondylotic patients and asymptomatic subjects. *Eur Spine J* 2015;24: 1265–73.
 12. Park MS, Kelly MP, Lee DH, et al. Sagittal alignment as a predictor of clinical adjacent segment pathology requiring surgery after anterior cervical arthrodesis. *Spine J* 2014; 14:1228–34.
 13. Lee S, Son E, Seo E. Factors determining cervical spine sagittal balance in asymptomatic adults: correlation with spinopelvic balance and thoracic inlet alignment. *Spine J V* 15 2015;705–12.
 14. Scheer JK, Tang JA, Smith JS, et al. Cervical spine alignment, sagittal deformity, and clinical implications. *J Neurosurg Spine* 2013; 19:141–59.
 15. Hirabayashi K, Miyakawa J, Satomi K, et al. Operative results and postoperative progression of ossification among patients with ossification of cervical posterior longitudinal ligament. *Spine*. 1981;6(4):354–64.
 16. Kurokawa T, Tsuyama N, Tanaka H, et al. Enlargement of the spinal canal by the sagittal splitting of the spinous processes (in Japanese). *BessatsuSeikeigeka*. 1982; 2:234–40.
 17. Okada M, Minamide A, Endo T, et al. A prospective randomized study of clinical outcomes in patients with cervical compressive myelopathy treated with open-door or French-door laminoplasty. *Spine*. 2009; 34:1119–26.
 18. QIU Sujun, SUN Yu, ZHANG Fengshan, et al. No effect of open-door laminoplasty together with preservation of the unilateral paraspinal muscle ligament complex on the posterior muscle volume of cervical spine. *Chinese J Spine Spinal Cord* 2010; 20:401–5.
 19. Tsuru M, Nagata K, Ueno T, et al. Electron microscopic observation of established chondrocytes derived from human intervertebral disc hernia (KTN-1) and role of macrophages in spontaneous regression of degenerated tissues. *Spine J* 2001; 1:422–31.
 20. Komori H1, Shinomiya K, Nakai O, et al. The natural history of herniated nucleus pulposus with radiculopathy. *Spine (Phila Pa 1976)* 1996; 21:225–9.
 21. Ahn SH, Ahn MW, Byun WM. Effect of the transligamentous extension of lumbar disc herniations on their regression and the clinical outcome of sciatica. *Spine (Phila Pa 1976)* 2000; 25:475–80.
 22. Buttermann GR. Lumbar disc herniation regression after successful epidural steroid injection. *J Spinal Disord Tech* 2002; 15:469–76.
 23. Haro H, Komori H, Kato T, et al. Experimental studies on the effects of recombinant human matrix metalloproteinases on herniated disc tissues: how to facilitate the natural resorption process of herniated discs. *J Orthop Res* 2005; 23:412–9.
 24. [24] Guinto FC, Hashim H, Stumer M. CT demonstration of disk regression after conservative therapy. *Am J Neuroradiol* 1984; 5:632–3.
 25. George J, Haskin E. Spontaneous regression of herniated nucleus pulposus. *AJR Am J Roentgenol* 1985; 145:371–5.
 26. Hirabayashi S1, Kumano K, Tsuiki T, et al. A dorsally displaced free fragment of lumbar disc herniation and its interesting histologic findings. *Spine (Phila Pa 1976)* 1990;15:1231–3.