

## Original Research Article

# MORPHOMETRIC ANALYSIS OF INDEX FINGER AND RING FINGER AND COMPARING WITH HEIGHT WEIGHT AND BMI

Dilip Kumar Sudele<sup>1</sup>, Heena Chaudhari<sup>2</sup>, Pankaj Singh<sup>3</sup>, Mridul Tripathi<sup>4</sup>

<sup>1</sup>Assistant Professor, Department of Anatomy, Raipur Institute of Medical Sciences, Bhansoj Road, Raipur, India.

<sup>2</sup>Associate Professor, Department of Anatomy, GMERS medical College, Valsad, India.

<sup>3</sup>Assistant Professor Department of Anatomy, Saraswathi Medical College, Unnao, UP, India.

<sup>4</sup>Assistant Professor, Department of Anatomy, Raipur Institute of Medical Sciences, Bhansoj Road, Raipur, India.

Received : 27/03/2024  
Received in revised form : 21/05/2024  
Accepted : 06/06/2024

### Corresponding Author:

**Dr. Mridul Tripathi**  
Assistant Professor, Department of Anatomy, Raipur Institute of Medical Sciences, Bhansoj Road, Raipur  
Email: drmridultripathi@yahoo.com

DOI: 10.5530/ijmedph.2024.2.128

Source of Support: Nil,  
Conflict of Interest: None declared

Int J Med Pub Health  
2024; 14 (2); 670-673

### ABSTRACT

**Background:** Sex determination in identification of human remains is one of the most important components in forensic identification and it is a major task in medico-legal investigation. **Aim:** Morphometric analysis of index finger and ring finger and comparing with height weight and BMI

**Material and Methods:** Present study was carried out on 400 individuals of both sexes. Length of index and ring fingers and the ratios between them were estimated.

**Results:** The mean length of Index finger of male was 7.049 cm and female 6.521 cm. The mean length of ring finger was in male and female were following 7.20 & 6.722 cm. The mean height of male was 170.7 and in female 158.5, weight of male 65.36 and female 48.41 where as both male and female BMI was 22.8 & 19.80.

**Conclusion:** Morphometric measurements of ring fingers and index finger and comparing it with height and weight of subject, the index finger and ring finger length ratio in males is greater than in females which is insignificant. In males' height was greater than the height of females. Weight and BMI of the males were greater than that of the females. In males out of all the anthropometric parameters, only BMI was significantly correlated with index finger length. In males no anthropometric parameters- was significantly correlated with ring finger length. In females neither index nor ring finger showed any significant co- relation with all the three anthropometric parameters.

**Keywords:** Human identification, Forensic science, Sex determination, Index/ring finger.

## INTRODUCTION

Anthropometry is a science which deals with method and techniques of measurement of living as well as skeletons of individuals.<sup>[1]</sup>

The evidence of use of this branch of science in the field of legal medicine is available since 19th century when a French police expert Mr. Alphonse Bertillon defined the system of criminal identification of individuals based on anthropometric measurements. Most of the workers used intact long bones like femur, tibia, humerus and radius for estimation of stature which is natural heights of a person in an

upright position and various methods, formulae and regression equations are devised for various population groups.<sup>[2]</sup>

Since anthropometric measurements are extensively used in the field of forensic and legal medicine. Estimation of stature is very important and main component of anthropological research which is necessary for identification and used by medico legal experts, forensic examiners and anatomist.<sup>[3]</sup>

If whole body is available then estimation of stature is easy but it is very difficult in case of only few parts of the body or some skeletal remains are available.<sup>[4]</sup>

Krogman WM in his research found the correlation between height and various finger lengths and also compared the stature with the finger length ratio.<sup>[5]</sup> Tyagi et al, found accurate or near accurate correlation between stature and finger lengths in Delhi region of India. Each race and population of geographical area needs specific formulae as it is because many factors like race, ethnicity, nutritional factors, and environmental status play important role in the development and growth of human beings.<sup>[6]</sup>

## MATERIAL AND METHODS

**Study Population:** - Study were carry out of 400 subject which was student of different batches of college, out of which 200 male and 200 female examined in Raipur institute of Medical Science (RIMS) they were of cosmopolitan origin ranging age (18-25) years. A written consent was obtained by each student in prescribed format.

### Instrument

1. Vernier caliper
2. Weight machine.



### Measurement of height



**Stature:** Maximum distance from vertex to floor, maintaining the anatomical

### Measurement of index and ring finger length

The measurements related to index & ring finger are taken with simple Vernier Caliper to measure the distance. It was measured straight distance from the midpoint of the proximal finger crease to the tip of the finger. Vernier caliper was used to measure the finger length; hand placed on the plane surface; palm of the hand is facing upwards. The dimension is taking three times by the same person and means are taking thus increasing the accuracy of the data. Where a band of crease the most proximal crease are used. The subject is properly sitting or standing in position. All fingers is fully extended palmar region with digit fully stretched touching a flat hand surface and 2<sup>nd</sup> and 4<sup>th</sup> digit adducted with the thumb slightly extended.

**Measuring of weight:** Subject was standing on weight machine without jacket and shoes.

### Inclusion Criteria

The criteria for including the individuals from the study was as follows-

1. Subject in age group of the 18-40 years.
2. Healthy individuals without any deformity or abnormality.

### Exclusion Criteria

1. Age group below 18 and above 40
2. Any type of physical deformity.
3. Any type of congenital anomalies.
4. Any of the finger injury.

**Statistical Analysis:** Statistical analysis was performed by using computer-based software, Statistical Package for Social Science (SPSS). Mean values of parameters were compared to determine

## RESULTS

According to the our present study the measurements of stature, weight, index and ring finger were selected by using direct physical procedure among 400 subjects (200 = Male and 200 = Female). The Mean and Standard Deviation were

calculated for each variable. Gender wise distribution of Height, Weight and BMI were studied. In our current study mean age of male subject were 18.94 and in female 19.34 it was show in table no 1. The mean length of Index finger of male was 7.049 cm and female 6.521 cm. the mean length of ring finger was in male and female following 7.20 & 6.722 cm data was given in table number 2. The mean height of male was 170.7 and in female 158.5, weight of male 65.36 and female 48.41 where as both male and female BMI was 22.8 & 19.80 show in table number 3.

**Table 1: Mean age distribution**

AGE DISTRIBUTION			
(GROUP)	(NUMBER)	(MEAN)	(SD)
MALE	200	18.94	1.49
FEMALE	200	19.34	1.59

**Table 2: Comparison of measurement of index & ring finger length in males & females**

S.NO	Male		Female		t-value	P -value
	Mean	SD	Mean	SD		
Index	7.049	0.470	6.521	0.504	7.65	<0.05*
Ringer	7.20	0.476	6.722	0.424	7.52	<0.05* (significant)
Index : Ring finger	0.97	0.0323	0.969	0.041	1.78	>0.05 (In significant)

**Table 3: General statistics of the anthropometric parameters used in males and female**

Parameter	Sex	Mean
Height	Male	170.7
	Female	158.585
Weight	Male	65.36
	Female	48.41
BMI	Male	22.8
	Female	19.8

## DISCUSSION

Determination of finger length ratio index finger and ring finger was done in cross section of sample of North India and the results were compared between males and females. According to previous researches done in the field shows that the digit ratio values are consistently reliable in determination of sexual dimorphism and finger length ratio in north Indian population.

Tyagi et al. observed that the positive correlation between stature and finger length and also suggested that the index finger was best for the prediction of stature in both the sexes studied in Delhi's population.<sup>[6]</sup> According to Bailey et al, Observed that males with lower finger length ratios had significantly higher physical aggression scores.<sup>[7]</sup>

Krishan et al found statistically significant correlation between stature, index finger length and ring length in right hand. Pearson correlation (r) was higher among males than females. Among males and females' correlation coefficient was higher for the index finger length than the ring finger length.<sup>[8]</sup> According to Brown et al, considerable proportion of normal males have low digit ratios compared to females.<sup>[9]</sup> Burley et al was also shown

that males had relatively short second finger than fourth finger.<sup>[10]</sup>

The present study was done to establish the relationship between the male and female finger length ratios index finger and ring finger equal and to ascertain if it has any correlation with height, weight and BMI. The result showed that there was correlation between index finger and ring finger and height in females of north Indian population and also observed that male had relatively short second digit than fourth digit.

The association of finger length ratio index finger and ring finger of Ebir ethnic group of nageria with physical traits was done and the results was compared between males and females of adult Ebir people. Many researchers have attempted these measurements but the digit ratio values proved to be consistently reliable in sex dimorphism and it has been demonstrated that a considerable proportion of normal males have low digit ratios compared to the females.<sup>[11]</sup>

The present study was designed to establish the relationship between the male and female finger length ratios index finger and ring finger mostly equal in north Indian population. In a study done by Manning et al show that the females have longer second digits than fourth digits while males have

longer fourth digits than second digits which was associated with height and weight. This accounted for the higher digit ratios in females than in males. The results of the males and females index finger and ring finger ratios however confirmed that digit ratios index finger and ring finger are sexually dimorphic phenomenon.<sup>[12]</sup>

Current study shows that the females have almost equal index finger and ring finger length and males also have equal length of both fingers incorporated with height and weight. Ratio of the digits in both sexes was also similar and it was statistically insignificant correlation between males and females.

## CONCLUSION

In male subjects we found that index finger is longer than the female subjects and the length of the ring finger was longer in males than the females. The index finger and ring finger length ratio in males is greater than in females which is insignificant. In males' height was greater than the height of females. Weight and BMI of the males were greater than that of the females. In males out of all the anthropometric parameters, only BMI was significantly correlated with index finger length. In males no anthropometric parameters- was significantly correlated with ring finger length. In females neither index nor ring finger showed any significant co- relation with all the three anthropometric parameters i.e. Height, Weight and BMI. In both males & females no statistically, significant difference was found in correlation

between ratio (2D:4D) with anthropometric parameters.

## REFERENCES

1. Singh IP, Bhasin MK. Anthropometry – a laboratory manual of biological anthropology. 1968, Delhi: kamal raj Enterprises 1-35.
2. Waghmare VKR, Gaikwad RB, Herekar NG. Estimation of the stature from the anthropometric measurement of hand length. The internet journal of biological Anthropology.2011 4(2).
3. Krishan k, kanchan t, dimaggio JA. A study of limb asymmetry & its effect on estimation of stature in forensic case work, Forensic SciInt 2010, 200(1-3).
4. Momonchand A, Devi Merra TH. Determination of stature from the interacromial length. JFMT 1999, 16(1) 72-73.
5. Krogman WM. The human skeleton in forensic medicine. 2ndspringfield: Charles C Thomas, 1986, 153-187.
6. Tyagi AK., Kohli A., Verma SK. Correlation between stature and finger length. International journal of medical Toxicology and legal medicine, 1999; 1(2): 20-22.
7. Bailey A, Hurd P. Finger length ratio (2D:4D) Correlates with psychology 2005; 68(3):215-222.
8. Krishan K, Kanchan T, Asha N. Estimation of stature from index finger and ring finger length in North Indian adolescents population. Journal of Forensic and legal Medicine, 2012; 19(5):285-290
9. Brown, W Hines, M, Fane B, Breeslove S. Masculinized finger length patterns in human males and females with congenital adrenal hyperplasia, Hormones and Behavior, 2002; 42(4), 380-86.
10. Burley N. T Faster VS. Digit ratio varies with sex, egg order and strength of mate preference in zebra finches. Proceedings of the Royal society B: Biological science 2004; 271,239-244.
11. Amitpratapdixit. Correlation between index finger and stature in uttrakhand population. 2014; 17(3): 1007-1009.
12. Manning J.T The ratio of 2nd to 4th digit ratio, sexual dimorphism, population differences and reproductive success: Evidence for sexually antagonistic genes?,Evol Hum Behav., 2000; 21(3): 163-183.