

Original Research Article

STUDY OF TREADMILL TEST IN DETECTING ASYMPTOMATIC CORONARY ARTERY DISEASE IN TYPE 2 DIABETES MELLITUS IN A TERTIARY CARE CENTER

M. Gayathri¹, J. Raj Kumar¹, K. Swetha¹

¹Assistant Professor, Department of General Medicine, Government Medical College, Krishnagiri, Tamil Nadu, India

Received : 09/10/2025
Received in revised form : 29/11/2025
Accepted : 13/12/2025

Corresponding Author:

Dr. M. Gayathri,
Assistant Professor, Department of
General Medicine, Government
Medical College, Krishnagiri, Tamil
Nadu, India.
Email: gayu3rockz@gmail.com

DOI: 10.70034/ijmedph.2025.4.559

Source of Support: Nil,
Conflict of Interest: None declared

Int J Med Pub Health
2025; 15 (4); 3120-3126

ABSTRACT

Background: Coronary Heart Disease is the common impact of death worldwide in type 2 Diabetes mellitus. High rate of asymptomatic coronary heart disease in type 2 diabetes mellitus. We studied about the prevalence of silent myocardial ischemia in asymptomatic type 2 diabetes mellitus by using treadmill test. We also analysed about the clinical predictions of silent myocardial ischemia in asymptomatic type 2 diabetes mellitus.

Materials and Methods: The study was conducted in the tertiary Care Hospital, Department of Medicine, during the period of July 2014 to July 2015. The study was one year cross sectional study conducted on the first 50 asymptomatic type 2 Diabetes mellitus outpatients without clinical and ECG evidence of CAD.

Results: Out of 50 cases, Treadmill test positive cases were 15 (30%) and negative cases were 35(70%) patients. 5 patients with DM duration between 16 and 20 years. Treadmill test was found to be positive in 4 (80%). 60% of subjects (8/13) with autonomic neuropathy had silent myocardial ischemia while only 18.9% of subjects (7/37) without autonomic neuropathy had myocardial ischemia. Average triglyceride in Treadmill test positive and negative cases 176.6 mg% and 130.5mg% respectively. Statistically significant value of $p < 0.001$ was found in triglycerides levels between both the groups. HbA1C (%) in positive and negative TMT cases were 9.7 and 7.7 respectively. Statistically significant value of $p < 0.001$ was found in HbA1C (%) levels between both the groups.

Conclusion: The preponderance of SMI in type 2 DM, without previous clinical and electrocardiographic signs of ischemic heart disease or hypertension is 30%. Age, duration of diabetes, triglyceride levels, autonomic neuropathy and glycosylated hemoglobin are strong clinical predictors of silent myocardial ischemia. Early detection of outpatients with type 2 DM for SMI may prevent catastrophic cardiac events. Based on our study, a routine screening is must for Silent Myocardial Ischemia in asymptomatic type 2 DM outpatients who had longer duration of diabetes, high triglycerides and HbA1C levels.

Keywords: Coronary Heart Disease, Diabetes Mellitus, Silent Myocardial Ischemia.

INTRODUCTION

Diabetes Mellitus (DM) is a disordered metabolic syndrome with hyperglycaemia either due to defect in insulin secretion or insulin resistance. It may cause progressive tissue damage and both micro and macrovascular complications. More than 95% of diabetes are type 2 DM. They always had insidious,

latent, along with asymptomatic phase. Therefore management of DM has changed not only controlling symptoms but also preventing complications.^[1-15]

BMI (Body Mass Index) of any indian of more than 23 must be investigated for diabetes.^[1] The waist circumference of more than 85cm in males and 80cm in females are the persons must be screened for DM.

The yearly check up for glycated haemoglobin and TOD (Target organ damage) is a must.^[16-22]

Coronary heart disease, atherosclerotic CHD, is the common impact of death worldwide in type 2 DM. INTERHEART -10 % of the population attributable risk of a first MI. High rate of asymptomatic coronary disease in type 2 DM. There will be a reduced myocardial flow reserve in type 2 DM.^[23-30]

Hyperglycaemia decreases endothelium-derived NO availability and affects vascular function mainly through the increased production of ROS. Asymptomatic DM higher median CAC scores raising frequency of silent ST segment depression and coronary perfusion changes during exercise testing.^[31-35]

Stress treadmill test is a readily available, cost effective, first line test for identification of coronary heart disease in DM of longer duration without any symptoms of angina.^[36-40]

This study was outlined to determine SMI (Silent Myocardial Ischemia) in selected asymptomatic DM outpatients by exercise tread mill test.^[2]

Aims & objectives

Aims:

1. To study the prevalence of silent myocardial ischemia in asymptomatic type 2 DM patients.

Objectives:

2. To study the prevalence of silent myocardial ischemia in asymptomatic type 2 DM patients by using tread mill test.
3. To analyse the clinical predictions of silent myocardial ischemia in asymptomatic type 2 DM patients.

MATERIALS AND METHODS

The study was conducted in the tertiary Care Hospital, Department of Medicine during the period of July 2014 to July 2015.

Study design: The study was one year cross sectional study on the asymptomatic type 2 DM outpatients without clinical and ECG evidence of CAD.

Source of data: The first 50 outpatients of asymptomatic type 2 DM without clinical evidence of CAD attending, diabetology and medicine OPD at tertiary care Hospital were enrolled in the study.

Sample size: A sample size of around 50 patients were included in the study.

Selection criteria:

1. All type 2 diabetic outpatients (both sex) in the age group of 35-60 yrs of age.
2. Asymptomatic outpatients with type 2 DM with normal resting ECG.

Exclusion criteria:

1. Previous history of MI, heart failure.
2. Evidence of angina pectoris
3. Anaemia
4. Hypertension
5. Renal disease
6. ECG evidence of Q wave MI, ischemic ST-segment or T wave abnormality or complete LBBB.
7. Any chronic illness because of cancer and ESRD or liver disease.

RESULTS

We studied around 50 outpatients of asymptomatic type 2 DM beyond clinical and electrocardiographic documentation of CAD in Coimbatore Medical college hospital, Coimbatore, and following results were noted.

Age: The mean age of study population was 48.62 yrs who were in the age group of 31-60 yrs of age. Most of the patients belonged to age group of 41-50 yrs of age.

Table 1: age and sex correlation

Age	Gender		Total	(%)
	Male	Female		
31-40	3	2	5	10%
41 – 50	22	4	26	52%
51 – 60	14	5	19	38%
Total	39	11	50	

In the age group of 31-40 yrs of age 5 patients are present. 3 were males, 2 were females. In the age group of 41-50 yrs of age 26 were present, 22 were

males, 4 were females. In the 51-60 yrs of age group 14 were males, 5 were females. Most patients in the age group of 41-50 yrs of age.

Mean Age with Gender						
Gender	Mean	SD	95% CI for Mean		Minimum	Maximum
	[Years]		Lower	Upper		
Male	48.59	6.07	46.62	50.56	35	60
Female	48.73	7.79	43.50	53.96	38	59
Total	48.62	6.40	46.80	50.44	35	60

Out of 50 cases 39 were males and 11 were females. In the age group of 31-40 yrs of age 5 patients are present. 3 were males, 2 were females. In the age group of 41-50 yrs of age 26 were present, 22 were

males, 4 were females. In the 51-60 yrs of age group 14 were males, 5 were females. Most patients in the age group of 41-50 yrs of age.

Table 2: Duration of diabetes mellitus

Duration of DM		
Duration	N	(%)
< 5 years	26	52%
6 – 10	12	24%
11 – 15	7	14%
> 15 years	5	10%
Total	50	100%

This table shows, more number of patients (26 i.e., 52%) were having diabetes equal to or less than 5 years, followed by 12 patients (24%) with the duration of 6-10 years, next 7 patients (14%) between 11-15 years and only 5 patients (10%) between 16-20 years.

In the study population, more number of patients (26 i.e., 52%) were having diabetes equal to or less than 5 years, followed by 12 patients (24%) with the duration of 6-10 years, next 7 patients (14%) between 11-15 years and only 5 patients (10%) between 16-20 years.

Table 3: glycosylated hemoglobin and duration of diabetes mellitus

Association of HbA1c with Duration of DM [N=50]						
HbA1c	Duration of DM				Total	(%)
	< 5 years	6 - 10	11 - 15	> 15 years		
6 – 8	21	1	2	0	24	48%
8 – 10	4	10	2	0	16	32%
10 – 12	1	1	3	5	10	20%
Total	26	12	7	5	50	

This table shows as there is increase in duration of DM, there is increase in HbA1C level. Out of 5 patients of more than 15 yrs duration of DM those glycemic index are very high around 10-12. It shows 5/5 (100%) of patients having poor glycemic index.

This shows as the duration of DM increased, there is a increase in HbA1C level. This shows poor glycaemic index. There is a statistically significant difference between duration of diabetes and HbA1C level.

Table 4: TMT results

TMT Results		
TMT	n	(%)
Positive	15	30%
Negative	35	70%
Total	50	100%

Out of 50 patients, Treadmill test was positive in 15 (30%) and Treadmill test was negative in 35 (70%) patients.

Table 5: TMT results and duration of diabetes mellitus

Duration	Positive	Negative	Total	(%)
< 5 years	3	22	26	52%
6 – 10	3	9	12	24%
11 – 15	5	2	7	14%
> 15 years	4	1	5	10%
Total	15	35	50	

Out of 50 cases Treadmill test was positive in 15 (30%) and negative in 35(70%) patients. Tread mill test was positive in 4/26 (27%), 3/9 (20%), 5/7 (33%) and 3/5(20%) patient with duration of diabetes ≤5, 6 to 10, 1 to 15, and 16-20 years respectively.

This shows the relation of silent myocardial ischemia and duration of diabetes mellitus. In the present study

out of 50 cases Treadmill test was positive in 15 (30%) and negative in 35(70%) patients. Tread mill test was positive in 4/26 (27%), 3/9 (20%), 5/7 (33%) and 3/5(20%) patient with duration of diabetes ≤5, 6 to 10, 1 to 15, and 16-20 years respectively. This shows longer the duration of diabetes, greater the risk of silent myocardial ischemia.

Table 6: comparison of diabetic subjects with and without silent myocardial ischaemia.

Mean of Clinical Variables with TMT Results							
Variables	TMT	Mean	SD	95% CI for Mean		Minimum	Maximum
				Lower	Upper		
Age	Positive	52.3	5.6	49.2	55.4	42	60
	Negative	47.1	6.2	44.9	49.2	35	60
	Total	48.6	6.4	46.8	50.4	35	60
Duration	Positive	10.4	5.3	7.5	13.3	2	18
	Negative	5.4	4.2	4.0	6.9	1	18
	Total	6.9	5.1	5.5	8.4	1	18

BMI	Positive	23.4	3.8	21.3	25.5	19.8	33.6	
	Negative	25.9	3.9	24.6	27.3	19.4	32.3	<0.05
	Total	25.2	4.0	24.0	26.3	19.4	33.6	
Pulse	Positive	75.9	6.1	72.6	79.3	68	86	
	Negative	76.0	6.0	73.9	78.1	68	90	>0.05
	Total	76.0	6.0	74.3	77.7	68	90	
SBP	Positive	127.9	3.2	126.1	129.6	120	130	
	Negative	128.4	4.7	126.8	130.0	120	136	>0.05
	Total	128.2	4.3	127.0	129.5	120	136	
DBP	Positive	74.3	4.8	71.6	76.9	70	80	
	Negative	75.3	5.4	73.5	77.2	60	84	>0.05
	Total	75.0	5.2	73.5	76.5	60	84	
FBS	Positive	170.5	33.1	152.2	188.9	126	256	
	Negative	183.0	40.3	169.1	196.8	125	291	>0.05
	Total	179.2	38.4	168.3	190.2	125	291	
PPBS	Positive	224.6	53.6	194.9	254.3	170	325	
	Negative	227.1	75.5	201.1	253.0	30	385	>0.05
	Total	226.3	69.1	206.7	246.0	30	385	
Cholesterol	Positive	174.4	31.8	156.8	192.0	133	234	
	Negative	180.9	28.5	171.2	190.7	133	234	>0.05
	Total	179.0	29.3	170.7	187.3	133	234	
TG	Positive	176.6	43.8	152.3	200.9	80	230	
	Negative	130.5	30.9	119.8	141.1	80	190	<0.001
	Total	144.3	40.8	132.7	155.9	80	230	
Urea	Positive	27.3	3.5	25.4	29.2	22	36	
	Negative	29.5	5.7	27.5	31.4	18	40	>0.05
	Total	28.8	5.2	27.3	30.3	18	40	
Creatinine	Positive	0.8	0.2	0.7	0.9	0.5	1.2	
	Negative	0.8	0.2	0.8	0.9	0.5	1.3	>0.05
	Total	0.8	0.2	0.8	0.9	0.5	1.3	
HbA1c	Positive	9.7	1.3	9.0	10.4	7.2	11.6	
	Negative	7.7	1.4	7.2	8.2	6.1	11.2	<0.001
	Total	8.3	1.6	7.8	8.8	6.1	11.6	

The observation made with treadmill testing is as follows with reference to average duration of diabetes, HbA1C levels and triglycerides levels; It was found that there was a statistically significant difference in Tread mill test positive cases and Tread mill test negative cases with reference to the average age, duration of DM ($p=0.001$), average HbA1C levels ($p=0.001$) and triglycerides levels ($p=0.001$). Average duration of DM in Treadmill test positive and negative cases was 10.4 and 5.4 years

respectively. Average age of diabetic patients in Treadmill test positive and negative cases are 52.3 and 47.1 respectively. Average HbA1C levels in Treadmill test positive and Treadmill test negative cases was 9.7% and 7.7% respectively. Average triglycerides levels in Treadmill test positive and negative cases was 176.6 and 130.5 mg% respectively.

Table 7: diabetic autonomic neuropathy and silent myocardial ischaemia.

Signs	TMT		
OH/ VM / SHG	Positive	Negative	Total
Yes	8	5	13
No	7	30	37
Total	15	35	50

8(61.5%) out of 13 DM with AN had silent myocardial ischemia, while 7(18.9%) out of 37 diabetics without autonomic neuropathy had silent myocardial ischemia.

8(61.5%) out of 13 DM with AN had silent myocardial ischemia, while 7(18.9%) out of 37 diabetes without AN had SMI. This shows DM with AN had high occurrence of SMI than without diabetes. There is a significant difference between autonomic neuropathy and SMI.

DISCUSSION

Coronary Artery Disease detection in asymptomatic type 2 DM is often delayed. The preponderance of

SMI in type 2 DM is variable and ranges from 9 to 75%.^[41-43] My study was anticipated at the asymptomatic presentation of CAD in the form of (SMI) in asymptomatic DM patients. It consisted of two aspects; First the prevalence of SMI in asymptomatic patients with type2 DM. Secondly to assess clinical predictors of SMI in these patients. This study consists of 50 known asymptomatic type 2 diabetics without clinical and electrocardiographic evidence of CAD and were evaluated for the preponderance of SMI by using exercise treadmill testing.

Out of 50 cases 39 were males and 11 were females. In the age group of 31-40 yrs of age 5 patients are present. 3 were males, 2 were females. In the age

group of 41-50 yrs of age 26 were present, 22 were males, 4 were females. In the 51-60 yrs of age group 14 were males, 5 were females. Most patients in the age group of 41-50 yrs of age.

25 (50%) patients were with normal body mass index, 16 patients (32%) were overweight while only 9 (18%) patients were obese.

In the study population, more number of patients (26 i.e., 52%) were having diabetes equal to or less than 5 years, followed by 12 patients (24%) with the duration of 6-10 years, next 7 patients (14%) between 11-15 years and only 5 patients (10%) between 16-20 years.

As the duration of diabetes increased there is a increase in HbA1C level in cases of duration of diabetes >15 yrs duration of diabetes. This shows poor glycemic index. There is a statistically significant difference between duration of DM and HbA1C level.

In our 50 patients, 2 (8%) was on diet control alone, 25 (50%) were on one or other OHA's, 11 (22%) were on one or other form of insulin while 12 (24%) were receiving both OHA's and insulin. There is a significant difference between duration of diabetes and treatment regimen.

Out of 50 cases TMT positive cases are 15 (30%) and negative cases are 35(70%) patients.

This study shows the prevalence of SMI in type 2 asymptomatic DM was turned out to be 30% (15/50). Our findings were similar to previous studies. One study,^[43] found that 29% diabetes who were asymptomatic for CAD had SMI on 24 hour ambulatory monitoring exercise electrocardiogram. Another similar study,^[44] had shown higher predominance of SMI in DM as compared to non diabetics.

Another study in India identified that 38.3% of DM beyond prior CAD had SMI on treadmill test.^[45,46]

Another study from India, reported 50% incidence of silent myocardial ischemia in diabetics on exercise electrocardiogram and 35% on ambulatory monitoring.^[47]

Yet another group found 12.1% of diabetics free of CAD to have SMI on exercise electrocardiogram testing.^[48]

Another study,^[49] found that SMI was positive in 14(46.7%) out of 30 DM patients by treadmill test.

One more study population identified that a total of 113/522 patients (22%) had SMI using stress testing in asymptomatic patients with type 2 diabetes mellitus.^[50]

One more study found that 62/500 patients (12.4%) had SMI in outpatients with type DM by using exercise electrocardiogram. So in my study it shows that DM patients had a higher prevalence of silent myocardial ischemia.^[51]

Duration of type 2 diabetes mellitus and silent myocardial ischemia.

In 50 outpatients, Treadmill test positive cases are 15 patients (30%) and negative cases are 35 patients (70%).

- 26 patients with diabetes duration equal to or less than 5 years, Tread mill test was positive in 3(11%).
- 12 patients with diabetes duration between 6-10years.Tread mill test was positive in 3(25%).
- 7 patients with DM duration between 11 and 15 years Treadmill test positive in 5(71.4%).
- 5 patients with DM duration between 16 and 20 years Treadmill test was found to be positive in 4 (80%).

Our results are similar to one study,^[52] that found that 70% subjects (7/10) with DM of <5 years duration had associated silent myocardial ischemia while only 30% subjects (3/10) with DM of >5years duration had associated SMI.

Another study,^[51] including 500 patients with type 2 diabetes mellitus with normal resting ECG found that, 62(12.4%) patients had asymptomatic coronary artery disease on exercise treadmill testing. The abnormalities of exercise test were associated with longer duration of diabetes($p<0.005$).

Silent myocardial ischemia and autonomic neuropathy: In our study, 60% of subject s (8/13) with autonomic neuropathy (AN) had silent myocardial ischemia while only 18.9% of subjects (7/37) without autonomic neuropathy had myocardial ischemia. Hence it depicts that, DM patients with AN had much higher occurrence of SMI than those without it (60%Vs18.9%) of studies.

One group,^[53] depicts that in a DM population of already known or doubtful CAD had SMI much more common with AN (92%Vs39%).

Another study in India⁵⁶ showed that 38.3% had SMI with a greater prevalence in those with AN (59%) than those without it (20%).

Another study,^[48] from India found that occurrence of SMI was higher in patients with AN 12/13(40%) compared to those without 3/30(10%) $p<0.001$.

Dyslipidemia and silent myocardial ischemia: In the present study, we found average total cholesterol in TMT positive and negative cases were 174.4mg% and 180.9mg % respectively. Average triglyceride in Treadmill test positive and negative cases 176.6mg% and 130.5mg% respectively. Statistically significant value of $p<0.001$ was found in triglycerides levels between both the groups.^[54]

The same results were observed in previous study which found that dyslipidemia was common in type 2DM and them most common aberrancy was elevated serum triglyceride levels (73.3%). The next aberrancy was decreased serum HDL levels(66.7%). Another study from India found that CAD had strong complementation with elevated levels of triglycerides (0.82) and low HDL (-0.81).^[55]

Yet another study found that triglyceride levels were elevated in 28 treadmill positive cases compared to 15 treadmill negative cases ($p<0.01$).

Glycosylated hemoglobin and silent myocardial ischemia.

The increased levels of glycosylated hemoglobin indicated poor glycemic control and it has great influence on CAD.

In our study we found average HbA1C (%) in Treadmill test positive and negative cases were 9.7 and 7.7 respectively. Statistically significant value of $p < 0.001$ was found in HbA1C (%) levels between both the groups.

One study found that among those who had diabetes mellitus, silent myocardial ischemia was present 27 of 54 patients (50%) who had hemoglobin A1c level $\geq 7.6\%$ and in 39 of 137 (28%) with hemoglobin A1c level ($p < 0.005$).

In one more study, it shows eloquent increasing trend of HbA1c levels over the augmenting number of coronary vessels involvement with CAD ($p < 0.0001$).

CONCLUSION

In our study population, the preponderance of SMI in type 2DM, without previous clinical and electrocardiographic signs of ischemic heart disease or hypertension is 30%. Duration of diabetes mellitus is directly proportional to increased risk of silent ischemia in type 2 DM. Triglyceride was found to be in higher levels in persons who had silent ischemia. Glycosylated hemoglobin levels was found to be more in diabetics, who had greater prevalence of silent myocardial ischemia on TMT.

Diabetics with clinical signs of autonomic neuropathy had higher incidence of silent myocardial ischemia. Age, duration of diabetes, triglyceride levels, autonomic neuropathy and glycosylated hemoglobin are strong clinical predictors of silent myocardial ischemia. Early detection of outpatients with type 2 DM for SMI may prevent catastrophic cardiac events. Based on our study, a routine screening for Silent Myocardial Ischemia in asymptomatic type 2 DM outpatients who had longer duration of diabetes, high triglycerides and HbA1C levels.

REFERENCES

1. American Diabetes Association-Clinical Practice Recommendations Diabetescare 2007;30(Suppl.4).
2. Fauci, Braunwald, Kasper, Hauser Longo, Jameson Loscalzo. Harrison's principles of internal medicine; Diabetes mellitus. 17th edn., New Delhi: McGraw Hill Medical Publishing Division, 2008.
3. API text book of medicine, 8th Edn. Association of physicians of India, Mumbai 2008.
4. Aiello LP. Diabetic retinopathy (Technical Review). Diabetes Care 1998;21:143-56.
5. UK Prospective Diabetes Study (UKPDS) Group: Effect of intensive blood glucose control with metformin on complications in overweight patients with type 2 diabetes (UKPDS 34). Lancet 1998; 352: 854-65.
6. Fuster V, Badimon L, Cohen M, et al: Insights into the pathogenesis of acute ischemic syndromes. Circulation 1988;77:1213-1220.
7. Fuster V, Badimon L, Badimon JJ, et al: The pathophysiology of coronary artery disease and the acute coronary syndromes. N Engl J Med 1992;326:242-250, 310-318.
8. Response to acetyl choline relates to risk factors for coronary artery disease. Circulation 1990;81:491-497.
9. Pasceri V, Willerson JT, Yeh EH: Direct pro-inflammatory effect of C-reactive protein on human endothelial cells. Circulation 2000;102:2165-2168.
10. Torzewski M, Rist C, Mortensen RF, et al: C-reactive protein: In the arterial intima: role of C-reactive protein receptor-dependent monocyte recruitment in atherosclerosis. Arterioscler Thromb Vasc Biol 2000; 20:2094-2099.
11. Yeh ET: CRP as a mediator of disease. Circulation 2004;109:II11-1114. Yeh ET, Anderson HV, Pasceri V, et al: C-reactive protein: linking inflammation to cardiovascular complications. Circulation 2001;104:974-975.
12. Blake GJ, Ridker PM: Novel clinical markers of vascular wall inflammation. Circulation Res 2001;89:763-771.
13. Liuzzo G, Biasucci LM, Gallimore JR, et al: The prognostic value of C-reactive protein and serum amyloid A protein in severe unstable angina. N Engl J Med 1994;331:417-424.
14. Kleinman J, Donahue R, Harris M, et al: Mortality among diabetics in a national sample. Am J Epidemiol 1988;128:389-401.
15. Butler W, Ostrander LJ, Carman W, Lamphiear D: Mortality from coronary heart disease in the Tecumseh study. Long-term effect of diabetes mellitus, glucose tolerance and other risk factors. Am J Epidemiol 1985;121:541-547.
16. Klein R: Hyperglycemia and microvascular and macrovascular disease in diabetes. Diabetes Care 1995;18:258-268.
17. Kuller LH, Velentgas P, Barzilay J, et al: Diabetes mellitus: Subclinical cardiovascular disease and all-cause mortality. Arterioscler Thromb Vasc Biol 2000;20:823-829.
18. The DECODE Study Group: Age, body mass index and glucose tolerance in 11 European population-based surveys. The DECODE Study Group, on behalf of the European Diabetes EPIDEMIOLOGY Group. Diabet Med 2002;19:558-565.
19. Lloyd CE, Kuller LH, Ellis D, et al: Coronary artery disease in IDDM: Gender differences in risk factors but not risk. Arterioscler Thromb Vasc Biol 1996;16:720-726.
20. Olson J, Edmundowicz D, Becker D, et al: Coronary calcium in adults with type 1 diabetes: A stronger correlate of clinical coronary artery disease in men than in women. Diabetes 2000;49:1571-1578.
21. Barzilay JI, Spiekerman CF, Kuller LH, et al: Prevalence of clinical and isolated subclinical cardiovascular disease in older adults with glucose disorders: The Cardiovascular Health Study. Diabetes Care 2001;24:1233-1239.
22. Pinhas-Hamiel O, Dolan LM, Dolan LM, Daniels SR, et al: Increased incidence of non-insulin-dependent diabetes mellitus among adolescents. J Pediatr 1996;128(5, Part 1):608-615.
23. Raheja BS. Heart disease in diabetes. Aetiopathogenesis in J of Assoc of Phy of Ind 1980; 28: 81-90.
24. Hume L, Oakley GD, Boulton AJ, Hardisty C, Ward JD. Asymptomatic myocardial ischemia in diabetes and its relationship to diabetic neuropathy. Diabetes Care 1986; 9: 389.
25. Achari V, Thakur AK. Treadmill testing in asymptomatic type 2 diabetes. JAPI 2002 Jan; 50: 50-53.
26. PK Garg, AK Swaroop, S. Singla. Silent myocardial ischaemia in type 2 diabetes mellitus. JAPI 2002 Dec; 50: 1471.
27. Koistinen MJ. Prevalence of asymptomatic myocardial ischemia in diabetic subjects. Br Med J 1990; 301: 92-5.
28. Marin Huerta E, Rayo I, Lara JI, Cuellar L, de la Calle H, Romero J et al. Silent myocardial ischemia during holter monitoring in patients with diabetes mellitus. Rev Esp Cardiol 1989; 42(8): 519-29.
29. Murray DP, O'Brien T, Mulrooney R, O'Sullivan. Autonomic dysfunction and silent myocardial ischemia on exercise testing in diabetes mellitus. Diabetic Med. 1990;8(7):580.
30. Scheidt-Nave C, Barrett-Connor E, Wingard DL. Resting ECG abnormalities suggestive of asymptomatic IHD associated with NIDDM in a defined population. Circulation 1990; 81: 899-906.
31. Turner RC, Millns H, Neil HAW, et al: Risk factors for coronary artery disease in non-insulin dependent diabetes mellitus: United Kingdom prospective diabetes study (UKPDS: 23) BMJ 1998;316:823-828.
32. Miettinen H, Lehto S, Salonen V, et al: Impact of diabetes on mortality after the first myocardial infarction. The FINMONICA Myocardial infarction register study group. Diabetes care 1998;21:69-75.

33. Zipes PD, Libby P, Bonow OR, Braunwald E. Braunwald's Heart Disease, 7th Edition, Philadelphia; WB Saunders company: 2005
34. Ellested, Myrvin H-Stress testing protocols, in stress testing principles and practice. 5th Edition, New Delhi: Jaypeeorthers; 2003
35. Ellested, Myrvin H-Stress testing protocols, in stress testing principles and practice. 5th Edition, New Delhi: Jaypeeorthers; 2003.
36. Janand-DelenneB,SavinB,Habib G, Bory M, Vague P, Lassmann-Vague V.Silent myocardial ischemia in patients with diabetes who to screen. Diabetes care 1999;22:1396-1400.
37. GorayaTY,leibsonCL, palumboPJ,etal.Coronary atherosclerosis in diabetes mellitus:a population-based autopsy study. J AM Collcardiol 2002;40:946-953.
38. KoistinenMJ.Prevalence of asymptomatic myocardial ischemia in diabetic subjects. BMJ 1990;301:92-95.
39. Scheidt-Nave C, Barrett-Connor E, Wingard DL. Resting ECG abnormalities suggestive of asymptomatic IHD associated with NIDDM in a defined population. Circulation 1990; 81: 899-906.
40. Motoji N. Silent myocardial ischemia in patients with NIDDM as judged by treadmill exercise testing and coronary angiography. Am Heart J 1991; 123:46.
41. Gupta SB, Pandit RB. Silent myocardial ischemia and cardiac autonomic neuropathy I diabetes. Ind Heart J 1993; 44(4): 227-9.
42. Ahluwalia G, Jain P, Chugh SK, Wasir HS, Kaul U. SMI in diabetics with normal autonomic function. Int J Cardiol 1995; 48(2): 147-53.
43. Misad Group. Prevalence of unrecognized SMI and its association with atherosclerotic risk factors in non insulin dependent DM. Am J Cardiol 1997; 79(2): 134-9.
44. Sukhija R, Chanwal D, Gambhir DS, Dewan R. Silent myocardial ischaemia in patients with type II diabetes mellitus and its relation with autonomic dysfunction. Indian Heart J 2000; 52(5): 540-6.
45. Wackers FJ, Young LH, Inzucchi SE, Chyun Da, Davey JA. Barrett EJ et al. Detection of Ischemia in Asymptomatic Diabetes Investigators. Detection of silent myocardial ischemia in asymptomatic diabetic subjects : the DIAD study. Diabetes Care 2004; 27(8): 1954-61.
46. Sargin H, Ozisik M, Oxisik NC, Seven O, Orbay E, Gozu H et al. The prevalence of silent ischemia in Turkish patients with type 2 diabetes mellitus. Tohoku J. Exp med. 2005; 20594: 351-5.
47. Ahuwalia V. Incidence of silent myocardial ischemic in non insulin dependent diabetes mellitus; 1998.
48. Murray DP, O'Brien T, Mulrooney R, O'Sullivan. Autonomic dysfunction and silent myocardial ischemia on exercise testing in diabetes mellitus. Diabetic Med. 1990;8(7):580
49. Langer A, Freeman MR, Josse RG, Steiner G, Armstrong PW. Detection of silent myocardial ischemia in DM. Am J Cardiol 1991; 67: 1073
50. Quek DK. Association of diabetic autonomic neuropathy and painless myocardial ischemia induced by exercise. Singapore Med J 1992; 32(2): 177-81.
51. Gupta SB, Pandit RB. Silent myocardial ischemia and cardiac autonomic neuropathy I diabetes. Ind Heart J 1993; 44(4): 227-9.
52. Mathura KC, Vaidya B, Gurbacharya DL. Study of serum lipid profile in type 2 diabetic patients attending KMCTH. Nepal Med Coll J 2005; 7(2): 97-100.
53. Agarwal R, Sharma P, Pal M, Kochar A, Kochar DK. Magnitude of dyslipidemia and its association with micro and macro vascular complications in type 2 diabetes; a hospital based study from Bikaner (Northwest India). Diabetes Res ClinPract 2006; 73(2): 211-4.
54. Achari V, Thakur AK. Treadmill testing in asymptomatic type 2 diabetes. JAPI 2002 Jan; 50: 50-53.
55. DeLuca AJ, Saulle LN, Aronow WS, Ravipathi G, Weis MB. Association of hemoglobin A1c with prevalence of silent myocardial ischaemia. Am J Cardiol 2005 June 15; 95(12): 1472-.
56. Gautam Ravipathi, Wilbert S. Aronow, Chulahn, et al. Association of hemoglobinA1c level with the severity of coronary artery disease in patients with diabetes mellitus. Am J Cardiol 2006 April 1; 97(7): 968- 969.